166 Pasadena Dr. Suite 100 Lexington, KY 40503

Office: 859-279-2111 Lexington
Plastic Surgery

Theodore L. Gerstle, M.D. www.lexingtonps.com drgerstle@lexingtonps.com

## ABDOMINOPLASTY

## REQUEST FOR TREATMENT AND INFORMED CONSENT

DO NOT SIGN THIS FORM UNTIL YOU HAVE READ IT AND FULLY U	NDERSTAND	ITS CC	NTENTS
PATIENT:	DATE:		/

I understand that the above named procedure has been explained and is to be performed on me.

The following has been explained to me in general terms and I understand that:

- The **DIAGNOSIS REQUIRING THIS PROCEDURE** is stretching and relaxation of the abdominal (belly) wall beyond normal, so that it protrudes and lacks support. This is most often the result of pregnancy or pregnancies, which have stretched the wall of the abdomen beyond its ability to return to it's normal supportive condition. Sometimes it is also related to a severe weight problem. This may affect the skin and muscles.
- 2. The **NATURE OF THE PROCEDURE** is to attempt to tighten the relaxed and stretched skin. This is called abdominoplasty and repair of rectus diastasis. This requires an incision across the low belly just above the hairline of the pubis from side to side and another incision around the umbilicus (belly button). The loose skin and fat can be pulled down and removed after it is tightened to reach the area of the low belly incision. The umbilicus is temporarily covered by upper abdominal skin when the belly wall is tightened downward. A new opening is then made and the umbilicus is brought out in its normal position. In some instances the umbilicus remains undisturbed or may be moved to a lower position. The skin repairs are made. Liposuction techniques which remove fat using a suction system may also be used as a component of the procedure. Areas of liposuction may include the belly and/or the hip and flank areas. Usually some temporary drains are placed to allow for tissue fluids to exit instead of causing swelling. Then the wounds are dressed.
- 3. The PURPOSE OF THIS PROCEDURE is to attempt to provide a tighter abdominal wall with more normal contour and better muscular support and relief of discomfort and problems, which are related. This procedure addresses excess skin and subcutaneous fat only.
- PRACTICAL ALTERNATIVES TO THIS PROCEDURE include doing nothing and accepting the circumstances of my medical 4. condition. Abdominal exercises or weight-loss may help some, but exercise may not return the stretched abdominal wall to normal.
- 5. IF I CHOOSE NOT TO HAVE THE ABOVE-NAMED PROCEDURE, MY PROGNOSIS (future medical condition) is not completely predictable and the medical condition may get better, may get worse or may stay the same. However, failure to have the procedure may result in possible progression of the medical condition and/or the possible need for more extensive surgery if the medical condition progresses and remains undiagnosed or untreated. Diet, exercise, pregnancy, aging and health problems may all contribute to future changes in my medical condition.
- 6. MATERIAL RISKS OF THIS PROCEDURE: As a result of this procedure being performed, there may be material risks of: INFECTION, ALLERGIC REACTION, TOXIC REACTION, DISFIGURING SCAR, SEVERE LOSS OF BLOOD, LOSS OR LOSS OF FUNCTION OF ANY LIMB OR ORGAN, BRAIN DAMAGE, CARDIAC ARREST OR DEATH
- 7. In addition to these material risks, there may be OTHER POSSIBLE RISKS involved in this procedure including but not limited to:

Initial \_\_\_\_ Page 1 of 3

166 Pasadena Dr. Suite 100 Lexington, KY 40503 Office: 859-279-2111

Theodore L. Gerstle, M.D. www.lexingtonps.com drgerstle@lexingtonps.com

- there will be permanent scars around the umbilicus and across the lower abdomen from side to side, above the hair line of the pubis; these scars generally go from pink and firm to faded and soft over a period of six to 12 months; some scars may widen, become depressed, or appear raised, firm and "ropey" red which may take two years or longer to fade and soften; some scars may pucker at the edges in an undesirable way; scars will be PERMANENT AND VISIBLE;
- 2) there may be an interference, temporary or permanent, with urination or bowel function;
- 3) bruising and swelling may occur and last a few weeks to several months;
- 4) contour irregularities or depressions may occur;
- 5) a hematoma (blood clot or collections of bloody fluid) may occur at the operative site;
- 6) severe blood loss may occur which may necessitate transfusion, which carries the risk of exposure to AIDS, hepatitis or other infectious diseases;
- 7) emboli or clots of blood and/or other material may go into the blood stream and travel to other parts of the body including the lungs or brain causing illness or even death;
- 8) abscess formation (collection of pus) may occur;
- 9) some tissue may slough (dissolve away) and result in a delayed healing situation and may require a skin graft (more common in smokers);
- 10) some fatty tissue may undergo fat necrosis (dissolve away) which may cause lumpiness or firmness in the tissue and may sometimes require drainage;
- 11) fluid collections may accumulate under the skin and may require drainage or aspiration (withdrawal by needle);
- 12) pain and discomfort may occur;
- 13) numbness (sensory loss, loss of feeling), itching, firmness, lumpiness and tight feelings may occur and could be temporary or permanent;
- 8. Even though the risks and complications cited above are infrequent, they are the ones peculiar to the operation and are of greatest concern. Complications may also be increased due to the patient's individual medical condition and personal habits. Medications, i.e. **ASPIRIN**, may interfere with blood clotting and cause excessive bleeding. **SMOKING CIGARETTES** may interfere with the blood supply to the skin and may cause abnormal healing with tissue sloughing (dissolving away) and excessive scarring. **ALCOHOL** may cause excessive bleeding during and after surgery. Certain **HERBAL PREPARATIONS** may affect the blood clotting system and cause excessive bleeding, while others may inhibit healing of the incisions. Colds, infections, boils and pustules may increase the risk of infection after surgery. Excessive sun exposure and/or tanning beds, heating pads and hot water bottles may cause severe burns at the surgery site if one has temporarily or permanently lost protective sensation.
- 9. I understand that the physician, medical personnel and other assistants will rely on statements made by me concerning my medical history and other information I provide in determining whether to perform the procedure or the course of treatment for my condition and in recommending the procedure which has been explained to me. Withholding medical and/or health information may result in further complications.
- 10. There may be a need for immediate or other additional surgery to treat the above complications, which could include hospitalization, time off work and additional expense to me.

Page 2 of 3 Initial \_\_\_\_\_

166 Pasadena Dr. Suite 100 Lexington, KY 40503

Office: 859-279-2111

Theodore L. Gerstle, M.D. www.lexingtonps.com drgerstle@lexingtonps.com

- 11. I understand that my expectations should be realistic and I should consider not undergoing the surgery if my expectations are greater than the reality of this treatment. Psychological problems may occur due to unrealistic expectations of the surgery or difficulties in accepting changes in my appearance.
- 12. I understand that the practice of medicine is not an exact science and that NO GUARANTEES OR ASSURANCES HAVE BEEN MADE TO ME CONCERNING THE RESULTS OF THIS PROCEDURE.
- 13. I consent to the taking of pictures during the course of my treatment for the purpose of helping to plan and assess the proposed therapy. No photographs will be shown to patients or physicians without my permission. If any portion of my surgery is to billed to insurance (this does not include cosmetic procedures), I understand my insurance carrier may require photographs to process my claim.
- 14. On occasion, surgical revisions may be indicated following the original surgery. If planned or performed within one (1) year after the original surgery, there will be no charge by the surgeon. However, a fee will be charged by the facility for use of the operating room. There will also be a charge by the anesthesiologist if indicated.
- 15. I voluntarily consent to allow Dr. Gerstle and all medical personnel under his direct supervision and control and all other personnel who may otherwise be involved in performing such procedures to perform the procedure(s) described or otherwise referred to herein.
- 16. BY SIGNING THIS FORM, I ACKNOWLEDGE THAT I HAVE READ OR HAD THIS FORM READ AND/OR EXPLAINED TO ME, THAT I FULLY UNDERSTAND ITS CONTENTS, THAT I HAVE BEEN GIVEN AMPLE OPPORTUNITY TO ASK QUESTIONS AND THAT ANY QUESTIONS HAVE BEEN ANSWERED SATISFACTORILY. ALL BLANKS OR STATEMENTS REQUIRING COMPLETION WERE FILLED IN.

Signature of person giving consent:	Date:
Relationship to patient if not the patient:	<del>-</del>
Witness:	Date:
Copy of consent form offered to patient	
Copy given Declined	